

Product Datasheet

Mono-Methyl-p53 (Lys370) (4C7) Mouse mAb

Catalog No: AB3631

Reactivity: Human, Mouse, Rat

Isotype: Mouse IgG1

Applications: IHC



www.abways.com

Information

UniProt ID: P04637

All Names: Cellular tumor antigen p53; Cys 51 stop; Tp53; Tumor protein p53

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide


Molecular Wt.: 53kDa

Application

IHC: 1/50-1/100

Background

Swiss-Prot Acc.P04637. Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkl1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2 (PubMed:24051492).



Immunohistochemistry analysis of paraffinembedded Human Lung Carcinoma using MonoMethylp53 (Lys370) (4C7) antibody.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com