

Product Datasheet

Smad3 (4H4) Mouse mAb

Catalog No: AB3639

Reactivity: Human, Mouse, Rat

Isotype: Mouse IgG1

Applications: WB, IHC



www.abways.com

Information

UniProt ID: P84022

All Names: SMAD3; MADH3; Mothers against decapentaplegic homolog 3; MAD homolog 3; Mad3; Mothers against DPP homolog 3; hMAD-3; JV15-2; SMAD family member 3; SMAD 3; Smad3; Hsmad3

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 50kDa

Application

WB: 1/500-1/1000

IHC: 1/50-1/100

Background

Swiss-Prot Acc.P84022.Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD3/SMAD4 complex, activates transcription. Also can form a SMAD3/SMAD4/JUN/FOS complex at the AP-1/SMAD site to regulate TGF-beta-mediated transcription. Has an inhibitory effect on wound healing probably by modulating both growth and migration of primary keratinocytes and by altering the TGF-mediated chemotaxis of monocytes. This effect on wound healing appears to be hormone-sensitive. Regulator of chondrogenesis and osteogenesis and inhibits early healing of bone fractures. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.



Western blot analysis of Smad3 in 293T, Jurkat, Rat Heart, Mouse Heart lysates using Smad3 (4H4) antibody.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com