

Product Datasheet

VEGF Receptor 1 Antibody

Catalog No: CY5126

Reactivity: Human, Mouse, Rat

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF IP FC



www.abways.com

Information

UniProt ID: P17948

All Names: VEGFR-1, VEGF Receptor 1, FLT-1, FLT, FRT, VEGFR1, Fms-like tyrosine kinase 1, Tyrosine-protein kinase FRT, Tyrosine-protein kinase receptor FLT

Form: Liquid

Storage instructions: Store at +4° C short term. Store at -20° C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: Synthesized peptide

Molecular Wt.: 151kDa

Application

WB: 1:500~1:2000


IHC: 1:50~1:200

ICC/IF: 1:50~1:200

IP: 1:50

Background

Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFB and PGF, and plays an essential role in the development of embryonic vasculature, the regulation of angiogenesis, cell survival, cell migration, macrophage function, chemotaxis, and cancer cell invasion. May play an essential role as a negative regulator of embryonic angiogenesis by inhibiting excessive proliferation of endothelial cells. Can promote endothelial cell proliferation, survival and angiogenesis in adulthood. Its function in promoting cell proliferation seems to be cell-type specific. Promotes PGF-mediated proliferation of endothelial cells, proliferation of some types of cancer cells, but does not promote proliferation of normal fibroblasts (in vitro). Has very high affinity for VEGFA and relatively low protein kinase activity; may function as a negative regulator of VEGFA signaling by limiting the amount of free VEGFA and preventing its binding to KDR. Likewise, isoforms lacking a transmembrane domain, such as isoform 2, isoform 3 and isoform 4, may function as decoy receptors for VEGFA.



Western blot analysis of VEGFR1 expression in A431 cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com