

Product Datasheet

Phospho-Smad3 (S423/S425) Antibody

Catalog No: CY5140

Reactivity: Human Mouse

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF



Information

UniProt ID: P84022

All Names: JV15-2, MAD-3, MADH3, Mad3, Mothers against DPP homolog 3, Mothers against decapentaplegic homolog 3, SMAD 3, Smad 3

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: Synthesized peptide

Molecular Wt.: 55kDa

Application


WB: 1:500~1:2000

IHC: 1:50~1:200

ICC/IF: 1:50~1:200

Background

Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD3/SMAD4 complex, activates transcription. Also can form a SMAD3/SMAD4/JUN/FOS complex at the AP-1/SMAD site to regulate TGF-beta-mediated transcription. Has an inhibitory effect on wound healing probably by modulating both growth and migration of primary keratinocytes and by altering the TGF-mediated chemotaxis of monocytes. This effect on wound healing appears to be hormone-sensitive. Regulator of chondrogenesis and osteogenesis and inhibits early healing of bone fractures. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.



Western blot analysis of Phospho-Smad3 (S423/S425) expression in A549 cell lysate treated with TGF- β 1.

For Research Use Only. Not For Use In Diagnostic Procedures.
www.abways.com