

Product Datasheet

CDK1 Antibody



Catalog No: CY5176

Reactivity: Human

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF IP

www.abways.com

Information

UniProt ID: P06493

All Names: CDC2; CDC28A; P34CDC2; MGC111195; DKFZp686L20222; CDK1

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 34 kDa

Application

WB: 1:500~1:2000


IHC: 1:50~1:200

ICC/IF: 1:50~1:200

IP: 1:50

Background

The cell division control protein cdc2, also known as cyclin-dependent kinase 1 (Cdk1) or p34/cdk1, plays a key role in the control of the eukaryotic cell cycle, where it is required for entry into S-phase and mitosis. Cdc2 exists as a complex with both cyclin A and cyclin B. The best characterized of these associations is the Cdc2 p34 cyclin B complex, which is required for the G2 to M phase transition. Activation of Cdc2 is controlled at several steps including cyclin binding and phosphorylation of threonine 161. However, the critical regulatory step in activating cdc2 during progression into mitosis appears to be dephosphorylation of Tyr15 and Tyr14. Phosphorylation at Tyr15 and inhibition of Cdc2 is carried out by WEE1 and MIK protein kinases while Tyr15 dephosphorylation and activation of Cdc2 is carried out by the cdc25 phosphatase. The isoform CDC2deltaT is found in breast cancer tissues. Furthermore, cdc2/Cdk1 is a key mediator of neuronal cell death in brain development and degeneration.



Western blot analysis of CDK1 expression in HeLa cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com

