

Product Datasheet

Bcl-6 Antibody



Catalog No: CY5184

Reactivity: Human

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF IP

www.abways.com

Information

UniProt ID: P41182

All Names: B-cell lymphoma 6 protein; BCL-6; B-cell lymphoma 5 protein; BCL-5; Protein LAZ-3; Zinc finger and BTB domain-containing protein 27; Zinc finger protein 51; BCL6; BCL5; LAZ3; ZBTB27; ZNF51

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 78 kDa

Application

WB: 1:500~1:2000


IHC: 1:50~1:200

ICC/IF: 1:50~1:200

IP: 1:50

Background

Bcl-6, a transcriptional repressor, binds Stat recognition-like DNA elements and influences germinal center development and cell differentiation. Additionally, Bcl-6 negatively regulates NFκB expression, thereby inhibiting NFκB-mediated cellular functions. HDAC- and silent information regulator (SIR)-2-dependent acetylation of Bcl-6 causes downregulation of activity by inhibiting the ability of Bcl-6 to recruit complexes containing histone deacetylases (HDACs). Bcl-6 is frequently deregulated in non-Hodgkin's B cell lymphomas. Bcl-6b (B-cell CLL/lymphoma 6, member B), also known as ZNF62, BAZF or ZBTB28, is a 480 amino acid nuclear protein that contains one BTB (POZ) domain and five C2H2-type zinc fingers. Expressed ubiquitously with highest expression in placenta and heart, Bcl-6b associates with Bcl-6 and functions as a sequence-specific transcriptional repressor that is thought to be necessary for early B-cell development.



Western blot analysis of Bcl-6 in expression Daudi cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com

