

Product Datasheet

Integrin beta 3 Antibody

Catalog No: CY5237

Reactivity: Human, Mouse ,Rat

Isotype: Rabbit IgG

Applications: WB, IHC, ICC/IF



www.abways.com

Information

UniProt ID: P05106

All Names: CD61; GP3A; GPIIIa; Integrin beta-3; integrin, beta 3 (platelet glycoprotein IIIa, antigen CD61); ITB3; ITGB3; platelet glycoprotein IIIa; Platelet membrane glycoprotein IIIa;

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 110 kDa

Application


WB: 1:1000~1:2000

IHC: 1:50~1:200

ICC/IF: 1:50~1:200

Background

Integrin alpha-V/beta-3 (ITGAV:ITGB3) is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin alpha-IIb/beta-3 (ITGA2B:ITGB3) is a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrins alpha-IIb/beta-3 and alpha-V/beta-3 recognize the sequence R-G-D in a wide array of ligands. Integrin alpha-IIb/beta-3 recognizes the sequence H-H-L-G-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin alpha-IIb/beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen. This step leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. Fibrinogen binding enhances SELP expression in activated platelets (By similarity). In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.



Western blot analysis of Integrin beta3 expression in human spleen lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com

