

Product Datasheet

PKC beta 2 Antibody

Catalog No: CY5390

Reactivity: Human Rat

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF IP FC



www.abways.com

Information

UniProt ID: P05771

All Names: Protein kinase C beta type; PRKCB; Prkcb; PKC-B; PKC-beta; PRKCB2; PKC Beta-II; EC:2.7.11.13

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 77 kDa

Application

WB: 1:500~1:2000


IHC: 1:50~1:200

ICC/IF: 1:50~1:200

IP: 1:50

Background

Calcium-activated, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase involved in various cellular processes such as regulation of the B-cell receptor (BCR) signalosome, oxidative stress-induced apoptosis, androgen receptor-dependent transcription regulation, insulin signaling and endothelial cells proliferation. Plays a key role in B-cell activation by regulating BCR-induced NF-kappa-B activation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by direct phosphorylation of CARD11/CARMA1 at 'Ser-559', 'Ser-644' and 'Ser-652'. Phosphorylation induces CARD11/CARMA1 association with lipid rafts and recruitment of the BCL10-MALT1 complex as well as MAP3K7/TAK1, which then activates IKK complex, resulting in nuclear translocation and activation of NFKB1.



Western blot analysis of PKC beta 2 expression in K562 cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com