

Product Datasheet

MyD88 Antibody

Catalog No: CY5681

Reactivity: Human

Isotype: Rabbit IgG

Applications: WB IHC ICC/IF FC



www.abways.com

Information

UniProt ID: Q99836

All Names: Myeloid differentiation primary response protein MyD88; MYD88;

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 33 kDa

Application

WB: 1:1000~1:2000


IHC: 1:50~1:200

ICC/IF: 1:50~1:200

FC: 1:100

Background

Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in *Drosophila*, play a pivotal role in innate immune responses (1-3). TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- κ B and subsequent regulation of immune and inflammatory genes. The TLRs and members of the IL-1 receptor family share a conserved stretch of approximately 200 amino acids known as the TIR domain. Upon activation, TLRs associate with a number of cytoplasmic adaptor proteins containing TIR domains including MyD88 (myeloid differentiation factor), MAL/TIRAP (MyD88-adaptor-like/TIR-associated protein), TRIF (Toll-receptor-associated activator of interferon), and TRAM (Toll-receptor-associated molecule). This association leads to the recruitment and activation of IRAK1 and IRAK4, which form a complex with TRAF6 to activate TAK1 and IKK. Activation of IKK leads to the degradation of I κ B that normally maintains NF- κ B inactivity by sequestering it in the cytoplasm.



Western blot analysis of MyD88 expression in Raji cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com