

Product Datasheet

SDF1 Antibody

Catalog No: CY5999

Reactivity: Human

Isotype: Rabbit IgG

Applications: WB ICC/IF



www.abways.com

Information

UniProt ID: P48061

All Names: Stromal cell-derived factor 1; SDF-1; hSDF-1; IRH; hIRH; PBSF; CXCL12; SDF1; SDF1A; SDF1B; Pre-B cell growth-stimulating factor; C-X-C motif chemokine 12;

Form: Liquid

Storage instructions: Store at +4° C short term. Store at -20° C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 11 kDa


Application

WB 1:1000~1:5000

ICC/IF 1:50~1:200

Background

Chemoattractant active on T-lymphocytes, monocytes, but not neutrophils. Activates the C-X-C chemokine receptor CXCR4 to induce a rapid and transient rise in the level of intracellular calcium ions and chemotaxis. Also binds to atypical chemokine receptor ACKR3, which activates the beta-arrestin pathway and acts as a scavenger receptor for SDF-1. SDF-1-beta(3-72) and SDF-1-alpha(3-67) show a reduced chemotactic activity. Binding to cell surface proteoglycans seems to inhibit formation of SDF-1-alpha(3-67) and thus to preserve activity on local sites. Acts as a positive regulator of monocyte migration and a negative regulator of monocyte adhesion via the LYN kinase.



Western blot analysis of SDF1 expression in Jurkat cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com